Dosing for Infants and Children

From Your Healthcare Professional





DOSE: Every 6-8 hours as needed. DO NOT GIVE MORE THAN 4 DOSES IN 24 HOURS.

Available in:



Infants' MOTRIN® **Concentrated Drops**

Active Ingredient:

Ibuprofen 50 mg (NSAID)* (in each 1.25 mL)

*Nonsteroidal



Children's MOTRIN® **Oral Suspension**

Active Ingredient: Ibuprofen 100 mg (NSAID)*

(in each 5 mL) *Nonsteroidal





mL = milliller		anti-inflammatory drug Berry		anti-inflammatory drug		Berry	
WEIGHT	AGE	Berry Flavor Use p	product only as directed.	Dye-Free Berry	Berry Flavor	Use product only as directed.	Dye-Free Berry
_	<6 MONTHS	_			_		
12-17 POUNDS	6-11 MONTHS	1.25 mL	1.25 mL 1.875	.625	_		
18-23 POUNDS	12-23 MONTHS	1.875 mL	11.25 mL 1875	625	_		
24-35 POUNDS	2-3 YEARS	_			5 mL	5 mL	
36-47 POUNDS	4-5 YEARS	_			7.5 m	L 7.5 mL	
48-59 POUNDS	6-8 YEARS	_			10 m	L 10 mL	
60-71 POUNDS	9-10 YEARS	_			12.5 ו	mL 12.5 mL	
72-95 POUNDS	11 YEARS	_			15 m	L 15 mL	

Kids' Wellness Tracker

This free, all-in-one tool from the makers of Children's TYLENOL® and Children's MOTRIN® makes it easy to track your child's height, weight, BMI, vaccines, symptoms, and medicines, plus calculate dosing.

Available on the App Store[™] and the Google Play Store $^{\text{\tiny{M}}}$





Google Play is a trademark of Google Inc. App Store is a service mark of Apple Inc. Standard download rates may apply.

IMPORTANT INSTRUCTIONS for Proper Use

Today's Date: _

This dosing recommendation from your doctor will expire in 14 DAYS.

- Read and follow the label on all MOTRIN® products.
- Repeat dose every 6-8 hours as needed.
- Do NOT give more than 4 doses in 24 hours.
- Use only the dosing device (syringe or dosing cup) that came with the product.